



Probing nuclear rates with Planck and BICEP2

E. Di Valentino, C. Gustavino, J. Lesgourgues, G. Mangano, A. Melchiorri, G. Miele, O. Pisanti

► To cite this version:

E. Di Valentino, C. Gustavino, J. Lesgourgues, G. Mangano, A. Melchiorri, et al.. Probing nuclear rates with Planck and BICEP2. *Physical Review D*, 2014, 90, pp.023543. hal-01120006

HAL Id: hal-01120006

<https://hal.science/hal-01120006>

Submitted on 24 Feb 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Probing nuclear rates with Planck and BICEP2

Eleonora Di Valentino,¹ Carlo Gustavino,² Julien Lesgourgues,³ Gianpiero Mangano,⁴ Alessandro Melchiorri,¹ Gennaro Miele,⁵ and Ofelia Pisanti⁵

¹*Physics Department and INFN, Università di Roma “La Sapienza”, Ple Aldo Moro 2, 00185, Rome, Italy*

²*INFN, Università di Roma “La Sapienza”, Ple Aldo Moro 2, 00185, Rome, Italy*

³*Institut de Théorie des Phénomènes Physiques, EPFL, CH-1015, Lausanne, Switzerland, and CERN, Theory Division, CH-1211 Geneva 23, Switzerland, and*

LAPTh (CNRS - Université de Savoie), BP 110, F-74941 Annecy-le-Vieux Cedex, France

⁴*INFN, Sezione di Napoli, Complesso Univ. Monte S. Angelo, Via Cintia, I-80126 Napoli, Italy*

⁵*Dipartimento di Fisica, Università di Napoli “Federico II” and INFN, Sezione di Napoli, Complesso Univ. Monte S. Angelo, Via Cintia, I-80126 Napoli, Italy*

Big Bang Nucleosynthesis (BBN) relates key cosmological parameters to the primordial abundance of light elements. In this paper, we point out that the recent observations of Cosmic Microwave Background anisotropies by the Planck satellite and by the BICEP2 experiment constrain these parameters with such a high level of accuracy that the primordial deuterium abundance can be inferred with remarkable precision. For a given cosmological model, one can obtain independent information on nuclear processes in the energy range relevant for BBN, which determine the eventual $^2\text{H}/\text{H}$ yield. In particular, assuming the standard cosmological model, we show that a combined analysis of Planck data and of recent deuterium abundance measurements in metal-poor damped Lyman-alpha systems provides independent information on the cross section of the radiative capture reaction $d(p, \gamma)^3\text{He}$ converting deuterium into helium. Interestingly, the result is higher than the values suggested by a fit of present experimental data in the BBN energy range (10 – 300 keV), whereas it is in better agreement with *ab initio* theoretical calculations, based on models for the nuclear electromagnetic current derived from realistic interactions. Due to the correlation between the rate of the above nuclear process and the effective number of neutrinos N_{eff} , the same analysis points out a $N_{\text{eff}} > 3$ as well. We show how this observation changes when assuming a non-minimal cosmological scenario. We conclude that further data on the $d(p, \gamma)^3\text{He}$ cross section in the few hundred keV range, that can be collected by experiments like LUNA, may either confirm the low value of this rate, or rather give some hint in favour of next-to-minimal cosmological scenarios.

PACS numbers: 98.80.Es, 98.80.Jk, 95.30.Sf

I. INTRODUCTION

Big Bang Nucleosynthesis (BBN, see e.g. [1] for a recent overview) offers one of the most powerful methods to test the validity of the cosmological model around the MeV energy scale. Two key cosmological parameters enter BBN computations, the energy density in baryons, $\Omega_b h^2$, and the effective neutrino number, N_{eff} , defined such that the energy density of relativistic particles at BBN is given by

$$\rho_{\text{rel}} = \rho_\gamma \left(1 + \frac{7}{8} \left(\frac{4}{11} \right)^{4/3} N_{\text{eff}} \right), \quad (1)$$

where ρ_γ is the Cosmic Microwave Background (CMB) photon energy density, given today by $\rho_{\gamma,0} \approx 4.8 \times 10^{-34} \text{ g cm}^{-3}$.

Recent measurements of CMB anisotropies obtained by the Planck satellite are in very good agreement with the theoretical predictions of the minimal ΛCDM cosmological model. They significantly reduce the uncertainty on the parameters of this model, and provide strong bounds on its possible extensions [2]. Assuming a given cosmological scenario and standard BBN dynamics, it

is now possible to infer indirectly from Planck data the abundance of primordial nuclides with exquisite precision. For example, assuming ΛCDM , the Planck constraint on the baryon density, $\Omega_b h^2 = 0.02207 \pm 0.00027$, can be translated into a prediction for the primordial deuterium fraction using the public BBN code `PARthENoPE` [3][33]

$$^2\text{H}/\text{H} = (2.65 \pm 0.07) \cdot 10^{-5} \quad (68\% \text{ C.L.}), \quad (2)$$

This constraint is competitive with the most recent and precise direct observations. Recently, the authors of Ref. [4] (see also [5]) presented a new analysis of all known deuterium absorption-line systems, including some new data from very metal-poor Lyman-alpha systems at redshift $z = 3.06726$ (visible in the spectrum of the quasar QSO SDSS J1358+6522) and at redshift $z = 3.04984$ (seen in QSO SDSS J1419+0829). Their result

$$^2\text{H}/\text{H} = (2.53 \pm 0.04) \cdot 10^{-5} \quad (68\% \text{ C.L.}), \quad (3)$$

is smaller than the (indirect, model-dependent) cosmological determination from CMB data, but with a comparable uncertainty.

These two deuterium abundance determinations, while broadly consistent, are off by about two standard deviations. This small tension might well be the result of small

experimental systematics, either in Planck or in astrophysical deuterium measurements. However, the point of this paper is to underline that current BBN calculations could also be plagued by systematics in the experimental determination of nuclear rates. As explained in the following, the main uncertainty for standard BBN calculations of ^2H comes from the rate of the radiative capture reaction $d(p, \gamma)^3\text{He}$. A recent review of the experimental status for this process can be found in [6]. The low energy limit of its cross section $\sigma(E)$ (or equivalently, of the corresponding astrophysical factor $S(E)$ [34]) is well-known thanks to the results of the underground experiment LUNA [7]. However, during BBN, the relevant energy range in the center of mass is rather around $E \simeq 30 - 300$ keV. For such energies, the uncertainty on the cross section is at the level of 6-10% when fitting $S(E)$ with a polynomial expression. This translates into a theoretical error on the primordial $^2\text{H}/\text{H}$ ratio of the order of 2% (for a fixed value of the baryon density and N_{eff}), comparable to the experimental error in the above cosmological determination (2) or astrophysical determination (3).

Recently, a reliable *ab initio* nuclear theory calculation of this cross section has been performed in [8–10]. The uncertainty on this prediction can be conservatively estimated to be also of the order of 7% [11]. However, the theoretical result is systematically larger than the best-fit value derived from the experimental data in the BBN energy range. By plugging the theoretical estimate of the cross section in a BBN code one finds that more deuterium is destroyed for the same value of the cosmological baryon density, and thus the predicted primordial ^2H abundance results to be smaller [11]. Interestingly, this could be a way to reconcile the slightly different values of $^2\text{H}/\text{H}$ measured in astrophysical data and predicted by Planck. Indeed, the result quoted in eq. (2) using the public BBN code **PARthENoPE** [3] relies on a value of the cross section $d(p, \gamma)^3\text{He}$ inferred from nuclear experimental data (the default value for the $d(p, \gamma)^3\text{He}$ rate used in the code was calculated in [12], and agrees at the 1.4% level with the best-fit result of [6]).

Further data on this crucial cross section in the relevant energy range might be expected from experiments such as LUNA. While waiting for such measurements one can find out to which extent the deuterium measurement of [4] can be made even more compatible with Planck predictions when the rate of the reaction $d(p, \gamma)^3\text{He}$ is treated as a free input parameter. We will address this issue assuming different cosmological models: the minimal ΛCDM model, ΛCDM plus extra radiation, a non spatially-flat universe, etc. This simple exercise points out that, remarkably, present CMB data are powerful enough to provide information on nuclear rates. Moreover, we will see that our results give independent support to the theoretical calculation of [10]. Of course, this close interplay between astrophysical observations and nuclear physics is not new. It is worth while recalling the role that the solar neutrino problem played in the quest

for a more accurate solar model, and the impact of this question on experimental efforts for measuring specific nuclear cross sections.

The paper is organized as follows. In the next section, we discuss in more details the nuclear rates which are most relevant for the determination of the primordial deuterium abundance and its theoretical error. We introduce a simplified way to parameterize the level of uncertainty still affecting the $d(p, \gamma)^3\text{He}$ reaction rate, found to be sufficient for our analysis. In Section III, we describe our method for fitting cosmological and astrophysical data. We present our results in Section IV, and discuss their implications in Section V.

II. THE PRIMORDIAL DEUTERIUM AS FUNCTION OF COSMOLOGICAL PARAMETERS AND NUCLEAR RATES

As well known, the theoretical value of the primordial $^2\text{H}/\text{H}$ abundance is a rapidly decreasing function of the baryon density parameter $\Omega_b h^2$. If we consider a slightly more general cosmological model with extra radiation, it grows as N_{eff} increases. Finally, this value depends on the cross section of a few leading nuclear processes, responsible for the initial deuterium production and its subsequent processing into $A = 3$ nuclei. More precisely, the calculation depends on the thermal rate of such processes, obtained by convolving their energy-dependent cross section $\sigma(E)$ with the thermal energy distribution of incoming nuclei during BBN. The four leading reactions are listed in Table I. Note that the uncertainties reported in the Table, like all other results quoted in this paper, unless otherwise stated, are calculated with a version of **PARthENoPE** where the $d(p, \gamma)^3\text{He}$ reaction rate is updated to the best fit determination of [6].

In the past, BBN calculations were based on the experimental determination of the cross section of nuclear processes, measured in laboratory experiments. The situation has changed recently, since detailed theoretical calculations are now available, at least for some reaction. For example, this is the case for the cross section of the neutron-proton fusion reaction $p(n, \gamma)^2\text{H}$, for which a very accurate result could be derived using pion-less effective field theory, with a theoretical error below the percent level [13, 14] (see e.g. [12] for further details). Using **PARthENoPE**, one can propagate this error to the primordial deuterium abundance. The resulting uncertainty is very small, $\sigma_{^2\text{H}/\text{H}} = 0.002 \cdot 10^{-5}$, i.e. of the order of 0.1% (for $\Omega_b h^2$ fixed at the Planck best-fit value).

The cross sections of $d-d$ fusion reactions, $d(d, n)^3\text{He}$ and $d(d, p)^3\text{H}$, are still determined using experimental data. They have been measured in the 100 keV range with a 1-2% uncertainty [15]. This leads to a propagated uncertainty on the deuterium primordial abundance at most of the order of 1%, see Table I.

The main source of uncertainty is presently due to the radiative capture process $d(p, \gamma)^3\text{He}$ converting deu-

Reaction	Rate Symbol	$\sigma_{2\text{H}/\text{H}} \cdot 10^5$
$p(n, \gamma)^2\text{H}$	R_1	± 0.002
$d(p, \gamma)^3\text{He}$	R_2	± 0.062
$d(d, n)^3\text{He}$	R_3	± 0.020
$d(d, p)^3\text{H}$	R_4	± 0.013

TABLE I: List of the leading reactions and corresponding rate symbols controlling the deuterium abundance after BBN. The last column shows the error on the ratio $^2\text{H}/\text{H}$ coming from experimental (or theoretical) uncertainties in the cross section of each reaction, for a fixed baryon density $\Omega_b h^2 = 0.02207$.

terium into helium. The present experimental status for the corresponding astrophysical factor $S(E)$ (where E is the center of mass energy) is reviewed in [6]. As we already mentioned, when fitting a polynomial expression for $S(E)$ to the raw data, now dominated by the LUNA results [7], one finds that the uncertainty at 68% C.L. grows from 6% in the low energy limit to 19% around 1 MeV. In the energy range relevant for BBN, the uncertainty is in the range 6-10%, which gives an error on the primordial deuterium abundance of order $\sigma_{2\text{H}/\text{H}} = 0.062 \cdot 10^{-5}$, as reported in Table I. This uncertainty is comparable to the experimental error estimated by [4], and dominates the error budget. In addition, the best fit value of $S(E)$ inferred from the data in the range $30 \text{ keV} \leq E \leq 300 \text{ keV}$ is lower than the theoretical result of [8, 10] by about 1σ . This difference may have an impact on the concordance of Planck results for the baryon density with the deuterium abundance measured by [4].

Using **ParthENoPE** with the best fit experimental cross section for the $d(p, \gamma)^3\text{He}$ reaction, one can check that the best fit value of the astrophysical determination of the deuterium abundance, $^2\text{H}/\text{H} = 2.53 \cdot 10^{-5}$ [4], corresponds to $\Omega_b h^2 = 0.02269$. However, in the case of the minimal cosmological model (i.e. the spatially flat ΛCDM model, with no extra relativistic species and $N_{\text{eff}} = 3.046$ [16]), we have seen that Planck data yield $\Omega_b h^2 = 0.02207 \pm 0.00027$ (68% C.L.). Hence there is a moderate 2σ tension, which could be relaxed either by assuming a more complicated cosmological model compatible with higher values of the baryon density, or by adopting the theoretical value of the $d(p, \gamma)^3\text{He}$ cross section [10]. In the latter case, if we stick to the ΛCDM model, the same range for the baryon density leads to

$$^2\text{H}/\text{H} = (2.58 \pm 0.07) \cdot 10^{-5}, \quad (4)$$

in nice agreement with the astrophysical determination at the 1σ level. In other words, increasing the $d(p, \gamma)^3\text{He}$ thermal rate has the same effect of increasing the cosmological baryon fraction.

This is illustrated in Fig. 1 where the likelihood function $L(\Omega_b h^2, R_2)$

$$L(\Omega_b h^2, R_2) = \exp \left(- \frac{(^2\text{H}/\text{H}_{th}(\Omega_b h^2, R_2) - ^2\text{H}/\text{H}_{ex})^2}{\sigma_{ex}^2} \right), \quad (5)$$




FIG. 1: The likelihood $L(\Omega_b h^2)$, assuming the astrophysical determination of the primordial deuterium abundance $^2\text{H}/\text{H}$ by Cooke et al. [4], adopting either the experimental best fit $R_2^{ex}(T)$ (solid) or *ab initio* calculation $R_2^{th}(T)$, (dashed) [10]. The star shows the Planck best fit value of $\Omega_b h^2$ in the minimal ΛCDM model.

is plotted versus baryon density in two different scenarios. Indices *th* and *ex* refer to the theoretical value of $^2\text{H}/\text{H}$ and to the experimental result of [4], respectively. The solid line corresponds to $R_2^{ex}(T)$ obtained by using the best fit of experimental values for the $d(p, \gamma)^3\text{He}$ cross section, while the dashed line relies on the theoretical prediction of the same cross section [10], whose corresponding rate is denoted by $R_2^{th}(T)$. The latter brings the agreement with the Planck ΛCDM value of $\Omega_b h^2$ from the 2σ to the 1σ level. Note that, in calculating those likelihoods, we only included the experimental error on astrophysical measurements of the deuterium fraction, $\sigma_{ex} = 0.05$. Indeed, our purpose is to show what the baryon probability could like after a future measurement campaign of the $d(p, \gamma)^3\text{He}$ astrophysical factor, assuming a small uncertainty and two different central values for this measurement. If the theoretical calculation of [10] was experimentally confirmed, the likelihood profile would shift to the dashed curve.

In the next section, we will generalize this study to non-minimal cosmological scenarios. The aim is to see whether, by combining CMB and BBN data, we can grasp some robust information on the value of the thermal rate R_2 preferred by cosmology. To this end, it is enough to parametrize the generic $R_2(T)$ in terms of an overall rescaling factor A_2 , namely $R_2(T) = A_2 R_2^{ex}(T)$, and use it in **ParthENoPE**. This approximation may sound too simplistic, but one can easily check that the ratio $R_2^{th}(T)/R_2^{ex}(T)$ is almost independent of temperature in the region relevant for BBN. For example using a constant rescaling factor $A_2 = 1.055$ one can mimic $R_2^{th}(T)$ with quite a good precision, and this conclusion holds for any value of $\Omega_b h^2$ in the range from 0.021 to 0.024, with at most a 0.2% difference in the predicted deuterium abundance. Hence, the use of a constant rescaling factor A_2 is reliable enough for our purpose, and offers the advantage of limiting the number of extra free parameters

to one.

Assuming this ansatz, we introduce the baryon likelihood function, $L(\Omega_b h^2, A_2)$, through

$$L(\Omega_b h^2, A_2) = \exp \left(- \frac{(^2\text{H}/\text{H}_{th}(\Omega_b h^2, A_2) - ^2\text{H}/\text{H}_{ex})^2}{\sigma_{ex}^2 + \sigma_{th}^2} \right), \quad (6)$$

where the theoretical value is a function of the baryon density and the $d(p, \gamma)^3\text{He}$ thermal rate rescaling factor A_2 , and again we use the experimental value and its squared uncertainty, see Eq. (3). Finally, σ_{th}^2 is the squared propagated error on deuterium yield due to the present experimental uncertainty on R_2 .

III. DATA ANALYSIS METHOD

Our main dataset consists in the Planck public data release of March 2013 [17], based on Planck temperature completed by WMAP9 polarization at low ℓ . We also consider the recent B modes polarization data (5 bins) from the BICEP2 experiment [18]. We combine these two CMB datasets (referred as Planck+WP and Planck+WP+BICEP2 respectively) with the deuterium abundance likelihood function $L(\Omega_b h^2, A_2)$ (referred as BBN).

Occasionally, we will also include the direct measurement of the Hubble constant by [19] (referred as HST), and information on Baryon Acoustic Oscillations by SDSS-DR7 at redshift $z = 0.35$ [20], by SDSS-DR9 at $z = 0.57$ [21], and by WiggleZ at $z = 0.44, 0.60, 0.73$ [22] (referred altogether as BAO).

For the data analysis method, we will use indifferently the publicly available Monte Carlo Markov Chain packages COSMOMC [23] (<http://cosmologist.info/cosmomc/>) and MONTE PYTHON [24] (<http://montepython.net>), which rely on the Metropolis-Hastings algorithm for exploring the parameter space, and on a convergence diagnostic based on the Gelman and Rubin statistics. We use the latest version of the two codes (April 201a), which include the support for the Planck Likelihood Code v1.0 (see <http://www.sciops.esa.int/wikiSI/planckpla/>) and implement an efficient sampling of the parameter space using a fast/slow parameter decorrelation [25]. We checked that the results from the two codes were identical. To evaluate the deuterium abundance produced during the Big Bang Nucleosynthesis, we use the **ParthENoPE** code, minimally modified in order to account for the global rescaling factor A_2 .

We will first consider the Planck+WP dataset assuming the minimal ΛCDM model with six free parameters: the density of baryons and cold dark matter $\Omega_b h^2$ and $\Omega_c h^2$, the ratio θ of the sound horizon to the angular diameter distance at decoupling, the optical depth to reionization τ , the amplitude A_S of the primordial scalar fluctuation spectrum at $k = 0.05 \text{ Mpc}^{-1}$, and the spectral index n_S of this spectrum. We extend this list of

free parameters to include the rescaling factor A_2 , affecting only the determination of the primordial deuterium abundance. For this model, we consider purely adiabatic initial conditions, we impose spatial flatness, we fix the effective number of neutrinos to its standard value $N_{\text{eff}} = 3.046$ [16], and we consider the sum of neutrino masses to be 0.06 eV as in the [2].

Subsequently, we will study several extensions of the minimal ΛCDM model, with extra free parameters: the neutrino effective number N_{eff} , the spatial curvature of the universe parametrised by $\Omega_k = 1 - \Omega_c - \Omega_b - \Omega_\Lambda$, and the amplitude of the lensing power spectrum A_L [26].

Finally, we consider a $\Lambda\text{CDM}+r$ framework where we allow the possibility for a gravitational wave background with tensor to scalar amplitude ratio r . In this case we include the BICEP2 dataset, assuming the B mode signal claimed by this experiment to be the genuine signature of primordial inflationary tensor modes. Since the amplitude of tensor modes measured by BICEP2 is in tension with the upper limit on r coming from the Planck experiment, we also consider two further extensions that could in principle solve the tension: an extra number of relativistic particles parametrized by N_{eff} (see e.g. [27]) and a running of the spectral index $dn_S/d\ln k$ [18].

IV. RESULTS

In Table II, we report our results for the parameters of the minimal ΛCDM model (plus the nuclear rate parameter A_2 and the derived cosmological parameter H_0), using the data combinations Planck+WP+BBN and PLANCK+WP+BBN+BAO.

As expected from the discussion of sections I and II, we find that the data provides an indication for A_2 being greater than one, roughly at the level of two standard deviations, even when adding the BAO dataset. We can also check explicitly in Figure 2 (top panel) that there is a clear anti-correlation between A_2 and $\Omega_b h^2$: in order to improve the agreement between Planck data and deuterium abundance measurements, one needs either a value of the nuclear rate rescaling factor A_2 higher than one, or a value of the baryon density larger than the Planck mean value. This is could be expected, since deuterium is a decreasing function of both the R_2 rate and the baryon density Ω_b . The lower panel of Figure 2 also shows an interesting correlation between A_2 and the Hubble constant H_0 . Letting A_2 vary yields a lower value for the Hubble constant in a combined Planck+WP+BBN analysis.

Given the fact that our results depend on the underlying cosmological model, it is interesting to investigate whether extensions of the standard ΛCDM model could bring the value of A_2 back in better agreement with the current experimental determination of $R_2(T)$ (corresponding by definition to $A_2 = 1$).

In Table III, we report the constraints when a variation in the neutrino effective number N_{eff} is allowed

Parameter	Planck+WP +BBN	Planck+WP +BBN+BAO
$\Omega_b h^2$	0.02202 ± 0.00028	0.02209 ± 0.00025
$\Omega_c h^2$	0.1200 ± 0.0026	0.1188 ± 0.0017
θ	1.04129 ± 0.00063	1.04144 ± 0.00058
τ	0.089 ± 0.013	0.091 ± 0.013
n_s	0.9599 ± 0.0073	0.9625 ± 0.0058
$\log[10^{10} A_s]$	3.089 ± 0.025	3.089 ± 0.025
$H_0[\text{km/s/Mpc}]$	67.2 ± 1.2	67.74 ± 0.78
A_2	1.155 ± 0.082	1.138 ± 0.076

TABLE II: Constraints on cosmological parameters (at the 68% confidence level) in the case of the minimal Λ CDM model.

Parameter	Planck+WP +BBN	Planck+WP +BBN+HST	Planck+WP +BBN+BAO
$\Omega_b h^2$	0.02241 ± 0.00042	0.02261 ± 0.00031	0.02233 ± 0.00029
$\Omega_c h^2$	0.1263 ± 0.0055	0.1281 ± 0.0049	0.1251 ± 0.0051
τ	0.096 ± 0.015	0.099 ± 0.014	0.094 ± 0.013
n_s	0.979 ± 0.017	0.988 ± 0.011	0.974 ± 0.010
$\log[10^{10} A_s]$	3.117 ± 0.034	3.128 ± 0.030	3.109 ± 0.029
$H_0[\text{km/s/Mpc}]$	71.0 ± 3.2	72.8 ± 2.0	70.1 ± 1.9
N_{eff}	3.56 ± 0.40	3.76 ± 0.27	3.43 ± 0.30
A_2	1.29 ± 0.15	1.33 ± 0.14	1.26 ± 0.14

TABLE III: Constraints on cosmological parameters (at the 68% confidence level) in the case of the extended Λ CDM model with extra relativistic degrees of freedom.

(to account, e.g., for extra relativistic degrees of freedom, or for non-standard physics in the neutrino sector). Even in that case, we can see that the combined Planck+WP+BBN and Planck+WP+BBN+BAO analyses show a preference for $A_2 > 1$ at roughly the 2σ level, even if the central value and error bar for A_2 are almost doubled. When the direct measurement of the Hubble parameter is included (case Planck+WP+BBN+HST), the indication for $A_2 > 1$ is even stronger, at the 2.5σ level. We can conclude that the preference for a large $d(p, \gamma)^3\text{He}$ reaction rate is robust against the extension of the minimal cosmological model to a free N_{eff} .

It is interesting to note that in Table III, the preferred value for the neutrino effective number N_{eff} is always larger than the standard value 3.046. As reported in section 6.4.4. of Ref. [2], the “standard” Planck+WP+BBN analysis (assuming $A_2 = 1$) gives $N_{\text{eff}} = 3.02 \pm 0.27$ (68% C.L.), while the CMB only result is $N_{\text{eff}} = 3.36 \pm 0.34$ (to be precise, in these results, the CMB dataset includes high- ℓ data from ACT and SPT, but the same trend is observed with only Planck+WP). With the present analysis, it becomes clear that this shift of N_{eff} towards its standard value is mostly driven by the low experimental value of R_2 . When A_2 is let free, the preference for $N_{\text{eff}} > 3.046$ persists even when deuterium measurements are included. This can also be checked in Fig. 3, where we report the two dimensional likelihood contours in the N_{eff} vs. A_2 plane for the three different datasets: Planck+WP+BBN, Planck+WP+BBN+HST, and Planck+WP+BBN+BAO. A correlation between A_2

and N_{eff} is clearly present: large values of A_2 remain compatible with Planck+WP+BBN data, provided that at the same time N_{eff} is larger than three. Such considerations reinforce the motivations for future experimental campaign to collect further data on the $d(p, \gamma)^3\text{He}$ cross section in the few hundred keV range. Notice that for $A_2 = 1.055$, corresponding to the theoretical result of [10] a standard value of N_{eff} is allowed at 68% C.L.. If experiments would confirm the theoretical result $R_2^{\text{th}}(T)$ in the BBN energy range, the overall agreement of CMB and BBN data for a standard number of relativistic degrees of freedom would improve with respect to the $A_2 = 1$ case. This does not hold if the HST measurement of H_0 is included in the analysis.

In Table IV we report the constraints on A_2 for further extensions of the minimal Λ CDM model, using the Planck+WP+BBN. We tried to vary the curvature parameter Ω_k , despite the fact that $\Omega_k \neq 0$ is difficult to explain from a theoretical point of view, and almost excluded when BAO data is also included. With free spatial curvature and without BAO data, the evidence for $A_2 > 1$ is slightly weaker. Finally, we considered the case of a free CMB lensing amplitude parameter A_L . Strictly speaking, this is not a physical extension of the Λ CDM model. The Planck data prefers $A_L > 1$, but as such, this result has no physical interpretation. It could be caused by a small and not yet identified systematic error affecting the Planck data (see the discussion in [2]), or alternatively, it may account in some approximate way for a non-standard growth rate of large scale structures

Parameter	Planck+WP+BBN	Planck+WP+BBN	Planck+WP+BBN	Planck+WP+BBN
$\Omega_b h^2$	0.02242 ± 0.00035	0.02301 ± 0.00051	0.02227 ± 0.00032	0.02261 ± 0.00042
$\Omega_c h^2$	0.1169 ± 0.0030	0.1245 ± 0.0055	0.1185 ± 0.0027	0.1241 ± 0.0053
θ	1.04179 ± 0.00067	1.04112 ± 0.00078	1.04153 ± 0.00065	1.04104 ± 0.00079
τ	0.087 ± 0.013	0.094 ± 0.015	0.087 ± 0.013	0.092 ± 0.015
n_s	0.9687 ± 0.0085	0.996 ± 0.018	0.9640 ± 0.0075	0.981 ± 0.015
$\log[10^{10} A_s]$	3.078 ± 0.025	3.111 ± 0.034	3.081 ± 0.025	3.105 ± 0.033
$H_0[\text{km/s/Mpc}]$	68.8 ± 1.4	74.3 ± 3.6	56.7 ± 5.4	5905 ± 6.4
N_{eff}	[3.046]	3.73 ± 0.40	[3.046]	3.50 ± 0.36
A_L	1.21 ± 0.12	1.25 ± 0.13	[1]	[1]
Ω_k	[0]	[0]	-0.035 ± 0.023	-0.035 ± 0.023
A_2	1.067 ± 0.086	1.21 ± 0.14	1.100 ± 0.084	1.21 ± 0.14

TABLE IV: Constraints on cosmological parameters (at the 68% confidence level) for several extensions of the Λ CDM model, with free parameters (N_{eff} , A_L , Ω_k). We vary at most two of these extra parameters at the same time, and fix the other ones to their standard model value, indicated above between squared brackets.

after recombination. We can see in Table IV that when A_L is left free, the A_2 parameter is well compatible with one. Our results for the joint confidence limits on A_2 vs. Ω_k and A_2 vs. A_L are shown in Fig.4.

In summary, Planck+WP+BBN data consistently indicate that $A_2 > 1$ (suggesting a $d(p, \gamma)^3\text{He}$ reaction rate closer to theoretical predictions than to experimental results) in the minimal Λ CDM model, as well as in a model with free N_{eff} . The evidence for $A_2 > 1$ goes away when either Ω_k or A_L are promoted as free parameters (with $N_{\text{eff}} = 3.046$), but these scenarios are less theoretically motivated. Incidentally, Table IV also shows that with a free Ω_k or A_L , and at the same time a free N_{eff} , the evidence for $A_2 > 1$ persists.

Finally, we have considered the Planck+WP+BICEP2+BBN dataset as stated in the previous section. In Table V we report the constraints using this dataset, allowing for a gravitational wave background with tensor to scalar ratio $r_{0.05}$ at scales of $k = 0.05 \text{ Mpc}^{-1}$. As we can see the indication for $A_2 > 1$ is still present in this case. Allowing for a variation in N_{eff} provides even further evidence for $A_2 > 1$ at more than two standard deviations. It is however interesting that when a running of the primordial spectral index is considered, A_2 is now compatible with one in between one standard deviation. In 5 we show the 2-D contour plots from the Planck+WP+BICEP2+BBN dataset in the $r_{0.05}$ vs A_2 (top panel), N_{eff} vs A_2 (center panel) and $dn_s/d\ln k$ vs A_2 (bottom panel) planes showing probabilities at 68% and 95%. As we can see, while there is essentially no degeneracy between A_2 and $r_{0.05}$, a degeneracy is clearly present between A_2 and N_{eff} and $dn_s/d\ln k$.

In summary, the BICEP2 dataset, when combined with the Planck data, provides an evidence either for a larger N_{eff} , either for a negative running of the spectral index $dn_s/d\ln k$. In the first case a value of A_2 strictly larger than one is needed in order to be in agreement with BBN. In the second case, when running is considered, A_2 is well compatible with one. A precise measurement of

A_2 from laboratory experiments could in principle help in a significative way in discriminating between these two scenarios.

V. CONCLUSIONS

In this work, we have shown that a combined analysis of Planck CMB data and of recent deuterium abundance measurements in metal-poor damped Lyman-alpha systems provides some piece of information on the radiative capture reaction $d(p, \gamma)^3\text{He}$, converting deuterium into helium. The value of the rate for this process represents the main source of uncertainty to date in the BBN computation of the primordial deuterium abundance within a given cosmological scenario, parameterized by the baryon density $\Omega_b h^2$ and effective neutrino number N_{eff} . The corresponding cross section has not been measured yet with a sufficiently low uncertainty and normalization errors in the BBN center of mass energy range, 30 - 300 keV. In addition to that, the best fit of available data appears to be systematically lower than the detailed theoretical calculation presented in [10]. Both these issues should be addressed by performing new dedicated experimental campaigns. We think that an experiment such as LUNA at the underground Gran Sasso Laboratories may give an answer to this problem in a reasonably short time.

In fact, with the present underground 400 kV LUNA accelerator [28] is possible to measure the $^2\text{H}(p, \gamma)^3\text{He}$ cross section in the $20 < E_{\text{cm}}(\text{keV}) < 260$ energy range with an accuracy better than 3%, i.e. considerably better than the 9% systematic uncertainty estimated in [29]. This goal can be achieved by using the large BGO detector already used in [30]. This detector ensures a detection efficiency of about 70% and a large angular coverage for the photons emitted by the $^2\text{H}(p, \gamma)^3\text{He}$ reaction. The accurate measurement of the $^2\text{H}(p, \gamma)^3\text{He}$ absolute cross section may be accomplished with the study of the angular distribution of emitted γ -rays by means of a large

Parameter	Planck+WP+BICEP2+BBN	Planck+WP+BICEP2+BBN	Planck+WP+BICEP2+BBN
$\Omega_b h^2$	0.02209 ± 0.00028	0.02286 ± 0.00044	0.02236 ± 0.00031
$\Omega_c h^2$	0.1184 ± 0.0027	0.1300 ± 0.0058	0.1195 ± 0.0027
θ	1.04146 ± 0.00063	1.04050 ± 0.00073	1.04144 ± 0.00063
τ	0.088 ± 0.012	0.100 ± 0.015	0.101 ± 0.015
n_s	0.9663 ± 0.0072	1.004 ± 0.018	0.9593 ± 0.0080
$\log[10^{10} A_s]$	3.082 ± 0.024	3.131 ± 0.034	3.115 ± 0.031
$H_0 [\text{km/s/Mpc}]$	67.9 ± 1.2	75.5 ± 3.7	67.7 ± 1.2
$r_{0.05}$	0.134 ± 0.045	0.153 ± 0.040	0.163 ± 0.040
N_{eff}	[3.046]	4.04 ± 0.44	[3.046]
$dn_s/d\ln k$	[0]	[0]	-0.0256 ± 0.0097
A_2	1.145 ± 0.081	1.40 ± 0.17	1.080 ± 0.079

TABLE V: Constraints on cosmological parameters (at the 68% confidence level) for the Planck+WP+BICEP2 dataset, with free parameters ($r_{0.05}, N_{\text{eff}}, dn_s/d\ln k$). We vary at most two of these extra parameters at the same time, and fix the other ones to their standard model value, indicated above between squared brackets.




FIG. 2: 2-D contour plots in the $\Omega_b h^2$ vs. A_2 (top panel) and H_0 vs. A_2 (bottom panel) planes, showing preferred parameter regions at the 68% and 95% confidence levels in the case of the minimal Λ CDM model.




FIG. 3: 2-D contour plots in the N_{eff} vs. A_2 plane, showing preferred parameter regions at the 68% and 95% confidence levels in the case of the extended Λ CDM model with extra relativistic degrees of freedom.

Ge(Li) detector [31, 32], in order to compare the data with "ab initio" modeling.

Our study shows that, interestingly, the combined analysis of Planck and deuterium abundance data returns a larger rate A_2 for this reaction than the best fit computed in [6], where the authors exploit the available experimental information on $d(p, \gamma)^3\text{He}$ cross section. On the other hand Planck is in better agreement with *ab initio* theoretical calculations. More precisely, when the reaction rate A_2 is chosen to match its present determination, Planck predicts a value of the primordial deuterium abundance in 2σ tension with its direct astrophysical determination. When the same reaction rate A_2 is assumed instead to match theoretical calculations, the two values of the primordial deuterium abundance agree at the 1σ level. We have shown that this conclusion holds in the minimal Λ CDM cosmological model, as well as when al-




FIG. 4: 2-D contour plots in the A_L vs A_2 (top panel) and Ω_k vs A_2 (bottom panel) planes showing probabilities at 68% and 95%.

lowing for a free effective neutrino number. In the latter case, the global likelihood analysis of astrophysical and cosmological data shows a direct correlation between A_2 and N_{eff} , so that higher values for A_2 are in better agreement with non standard scenarios with extra relativistic degrees of freedom.

Finally, we have shown that the inclusion of the new BICEP2 dataset also points towards a larger value for A_2 , especially when N_{eff} is left free to vary. However, a running of the spectral index could bring the value of A_2 back in agreement with one even when the BICEP2 dataset is considered.

New experimental data on the $d(p, \gamma)^3\text{He}$ reaction rate will therefore have a significant impact on the knowledge of N_{eff} and of $dn_s/d\ln k$ as well.

Acknowledgements

We are pleased to thank L. E. Marcucci, who kindly provided the results of the *ab initio* calculations of the $d(p, \gamma)^3\text{He}$ astrophysical factor described in [10] for discussions and help.

We thank the Planck Editorial Board for the internal refereeing of this work.

-
- [1] F. Iocco, G. Mangano, G. Miele, O. Pisanti and P. D. Serpico, Phys. Rept. **472**, 1 (2009) [arXiv:0809.0631 [astro-ph]].
 - [2] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5076 [astro-ph.CO].
 - [3] O. Pisanti, A. Cirillo, S. Esposito, F. Iocco, G. Mangano, G. Miele and P. D. Serpico, Comput. Phys. Commun. **178**, 956 (2008) [arXiv:0705.0290 [astro-ph]].
 - [4] R. Cooke, M. Pettini, R. A. Jorgenson, M. T. Murphy and C. C. Steidel, arXiv:1308.3240 [astro-ph.CO].
 - [5] M. Pettini and R. Cooke, Mon. Not. Roy. Astron. Soc. **425**, 2477 (2012) [arXiv:1205.3785 [astro-ph.CO]].
 - [6] E. G. Adelberger, A. B. Balantekin, D. Bemmerer, C. A. Bertulani, J. -W. Chen, H. Costantini, M. Couder and R. Cyburt *et al.*, Rev. Mod. Phys. **83**, 195 (2011) [arXiv:1004.2318 [nucl-ex]].
 - [7] C. Casella *et al.* [LUNA Collaboration] Nucl. Phys. A **706**, 203 (2002).
 - [8] M. Viviani, A. Kievsky, L. E. Marcucci, S. Rosati and R. Schiavilla, Phys. Rev. C **61**, 064001 (2000) [nucl-th/9911051].
 - [9] L. E. Marcucci, K. M. Nollett, R. Schiavilla and R. B. Wiringa, Nucl. Phys. A **777**, 111 (2006) [nucl-th/0402078].
 - [10] L. E. Marcucci, M. Viviani, R. Schiavilla, A. Kievsky and S. Rosati, Phys. Rev. C **72**, 014001 (2005) [nucl-th/0502048].
 - [11] K. M. Nollett and G. P. Holder, arXiv:1112.2683 [astro-ph.CO].
 - [12] P. D. Serpico, S. Esposito, F. Iocco, G. Mangano, G. Miele and O. Pisanti, JCAP **0412**, 010 (2004) [astro-ph/0408076].
 - [13] G. Rupak, Nucl. Phys. A **678**, 409 (2000).
 - [14] J.W. Chen and M.J. Savage, Phys. Rev. C **60**, 60205 (1999).
 - [15] D.S. Leonard *et al.*, Phys. Rev. C **73**, 045801 (2006) [nucl-ex/0601035].
 - [16] G. Mangano, G. Miele, S. Pastor, T. Pinto, O. Pisanti and P. D. Serpico, Nucl. Phys. B **729**, 221 (2005) [hep-ph/0506164].
 - [17] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5075 [astro-ph.CO].

- [18] P. A. R. Ade *et al.* [BICEP2 Collaboration], arXiv:1403.3985 [astro-ph.CO].
- [19] A. G. Riess, L. Macri, S. Casertano, H. Lampeitl, H. C. Ferguson, A. V. Filippenko, S. W. Jha and W. Li *et al.*, *Astrophys. J.* **730** (2011) 119 [Erratum-ibid. **732** (2011) 129] [arXiv:1103.2976 [astro-ph.CO]].
- [20] N. Padmanabhan, X. Xu, D. J. Eisenstein, R. Scalzo, A. J. Cuesta, K. T. Mehta and E. Kazin, *Mon. Not. Roy. Astron. Soc.* **427** (2012) 3, 2132 [arXiv:1202.0090 [astro-ph.CO]].
- [21] L. Anderson, E. Aubourg, S. Bailey, D. Bizyaev, M. Blanton, A. S. Bolton, J. Brinkmann and J. R. Brownstein *et al.*, *Mon. Not. Roy. Astron. Soc.* **427** (2013) 4, 3435 [arXiv:1203.6594 [astro-ph.CO]].
- [22] C. Blake, T. Davis, G. Poole, D. Parkinson, S. Brough, M. Colless, C. Contreras and W. Couch *et al.*, *Mon. Not. Roy. Astron. Soc.* **415** (2011) 2892 [arXiv:1105.2862 [astro-ph.CO]].
- [23] A. Lewis and S. Bridle, *Phys. Rev. D* **66**, 103511 (2002) [astro-ph/0205436].
- [24] B. Audren, J. Lesgourgues, K. Benabed and S. Prunet, *JCAP* **1302**, 001 (2013) [arXiv:1210.7183 [astro-ph.CO]].
- [25] A. Lewis, arXiv:1304.4473 [astro-ph.CO].
- [26] E. Calabrese, A. Slosar, A. Melchiorri, G. F. Smoot and O. Zahn, *Phys. Rev. D* **77**, 123531 (2008) [arXiv:0803.2309 [astro-ph]].
- [27] E. Giusarma, E. Di Valentino, M. Lattanzi, A. Melchiorri and O. Mena, arXiv:1403.4852 [astro-ph.CO]; J. -F. Zhang, Y. -H. Li and X. Zhang, arXiv:1403.7028 [astro-ph.CO]; C. Dvorkin, M. Wyman, D. H. Rudd and W. Hu, arXiv:1403.8049 [astro-ph.CO].
- [28] A. Formicola *et al.* (LUNA collaboration), *Nucl. Instr. and Meth. A* **507** (2003) 609.
- [29] L. Ma *et al.*, *Phys. Rev. C* **55**, 588 (1997).
- [30] C. Casella *et al.* (LUNA collaboration): *Nuclear Physics A* **706** (2002) 203216.
- [31] M. Anders *et al.* (LUNA collaboration): *Eur. Phys. J. A* (2013) **49**: 28.
- [32] M. Anders *et al.* (LUNA collaboration): *Phys. Rev. Lett.*, submitted.
- [33] In this paper we use a version of **ParthENoPE** where the $d(p, \gamma)^3\text{He}$ reaction rate is updated to the best fit experimental determination (see section II). The deuterium fraction given by the public version of **ParthENoPE** is slightly different, but the change in the central value is at the level of 4 per mille, only.
- [34] We recall that the energy-dependent cross section $\sigma(E)$ is related to the energy-dependent astrophysical factor $S(E)$ through $\sigma(E) = S(E)e^{-2\pi\eta}/E$, where η is the Sommerfeld factor.




FIG. 5: 2-D contour plots from the Planck+WP+BICEP2+BBN dataset in the r vs A_2 (top panel), N_{eff} vs A_2 (center panel) and $dn_s/d\ln k$ vs A_2 (bottom panel) planes showing probabilities at 68% and 95%.